

Can TeVeS avoid Dark Matter on galactic scales?

Nick E. Mavromatos,^{*} Mairi Sakellariadou,[†] and Muhammad Furqaan Yusa[‡]
King's College London, Department of Physics, Strand WC2R 2LS, London, U.K.

A fully relativistic analysis of gravitational lensing in TeVeS is presented. By estimating the lensing masses for a set of six lenses from the CASTLES database, and then comparing them to the stellar mass, the deficit between the two is obtained and analysed. Considering a parametrised range for the TeVeS function $\mu(y)$, which controls the strength of the modification to gravity, it is found that on galactic scales TeVeS requires additional dark matter with the commonly used $\mu(y)$. A soft dependence of the results on the cosmological framework and the TeVeS free parameters is discussed. For one particular form of $\mu(y)$, TeVeS is found to require very little dark matter. This choice is however ruled out by rotation curve data. The inability to simultaneously fit lensing and rotation curves for a single form of $\mu(y)$ is a challenge to a *no dark matter* TeVeS proposal.

PACS numbers:

The standard (Λ CDM) cosmological paradigm is based upon Cold Dark Matter (CDM), a cosmological constant Λ and classical general relativity/Friedman-Robertson-Lemaître-Walker cosmology. Despite its enormous success and consistency with a plethora of astrophysical data, competing models have been proposed for the primary reason of the still unknown nature of the dark energy component and the current undetectability of dark matter. To explain the observed flat rotation curves of galaxies without Dark Matter, Milgrom [1] proposed Modified Newtonian Dynamics (MOND), based upon the relation $f(|\vec{a}|/a_0)\vec{a} = -\vec{\nabla}\Phi_N$ between the acceleration $\vec{a}$ and the Newtonian gravitational field Φ_N . The constant $a_0 \approx 1.2 \times 10^{-10} \text{m/s}^2$ is motivated by the acceleration found in the outer regions of galaxies where the rotation curve is flat. When f , assumed to be a positive, smooth, monotonic function, equals unity, usual Newtonian dynamics hold, while when it approximately equals its argument, the deep MONDian regime applies.

MOND has been successful in explaining the dynamics of disk galaxies, however it is less successful for clusters of galaxies. It was promoted [2] to a classical relativistic field theory by introducing a TEnsor, VEcTOr and Scalar field (TeVeS). TeVeS has been criticised as lacking a fundamental theoretical motivation. Recently, it has been argued [3] that such a theory can emerge naturally within some string theory models.

In Ref. [4], where the lensing mass in MOND was compared to the stellar mass content of the lenses, it was found that comparable amounts of dark matter were needed in MOND to that required in the standard lensing scenario. This result is in contrast with attempts to explain the lensing data on galactic-cluster scales by introducing a 2 eV neutrino. In fact, this component of dark matter has been shown [5] to cluster on Mpc scales but not on galactic scales where the previous analysis was conducted. It was concluded that either lensing must operate in a qualitatively different way within the covariant “parent” theory of MOND, such as the TeVeS models, or dark matter should be considered within MOND even on

galactic scales.

In this letter TeVeS is examined in a similar way, namely by deriving the modified lensing equation and solving it numerically, to investigate whether it also shows a dark component from lensing. The (weak) dependence of the results on the cosmological models and TeVeS free parameters is then discussed. TeVeS [2] is a bi-metric model in which matter and radiation does not feel the Einstein metric, $g_{\alpha\beta}$, appearing in the canonical kinetic term in the (effective) action, but a modified “physical” metric, $\tilde{g}_{\alpha\beta}$, related to the Einstein metric by $\tilde{g}_{\alpha\beta} = e^{-2\phi}g_{\alpha\beta} - U_\alpha U_\beta (e^{2\phi} - e^{-2\phi})$, where U_μ, ϕ denote the TeVeS vector and scalar field, respectively. The TeVeS action is:

$$S = \int d^4x \left[\frac{1}{16\pi G} (R - 2\Lambda) - \frac{1}{2} \{ \sigma^2 (g^{\mu\nu} - U^\mu U^\nu) \phi_{,\alpha} \phi_{,\beta} + \frac{1}{2} G l^{-2} \sigma^4 F(kG\sigma^2) \} - \frac{1}{32\pi G} \{ K \mathcal{F}^{\alpha\beta} \mathcal{F}_{\alpha\beta} - 2\lambda (U_\mu U^\mu + 1) \} \right] (-g)^{1/2} + \mathcal{L}(\tilde{g}_{\mu\nu}, f^\alpha, f^\alpha_{|\mu}, \dots) (-\tilde{g})^{1/2}, \quad (1)$$

where k, K are the coupling constants for the scalar, vector field, respectively; ℓ is a free scale length related to a_0 (c.f below, after Eq.(5)); σ is an additional non-dynamical scalar field; $\mathcal{F}_{\mu\nu} \equiv U_{\mu,\nu} - U_{\nu,\mu}$; λ is a Lagrange multiplier implementing the constraint $g^{\alpha\beta} U_\alpha U_\beta = -1$, which is completely fixed by variation of the action; the function $F(kG\sigma)$ is chosen to give the correct non-relativistic MONDian limit, with G related to the Newtonian gravitational constant, G_N , by $G = G_N / (1 + \frac{K+k}{2} - 2\phi_c)$, where ϕ_c is the present day cosmological value of the scalar field. Covariant derivatives denoted by $_{|\mu} \dots$ are taken with respect to $\tilde{g}_{\mu\nu}$ and indices are raised/lowered with the metric $g_{\mu\nu}$. A new function $\mu(y)$ is introduced as [2]

$$-\mu F(\mu) - \frac{1}{2} \mu^2 \frac{dF(\mu)}{d\mu} = y \equiv k\ell^2 (g^{\mu\nu} - U^\mu U^\nu) \phi_{,\mu} \phi_{,\nu} \quad kG\sigma^2 = \mu (k\ell^2 (g^{\mu\nu} - U^\mu U^\nu) \phi_{,\mu} \phi_{,\nu}). \quad (2)$$

Possible choices for the μ function will be discussed later.

The isotropic spherically symmetric Einstein metric can be generically written as $g_{\alpha\beta}dx^\alpha dx^\beta = -e^\nu dt^2 + e^\zeta(dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2)$; both ν and ζ are functions of r . The physical metric $\tilde{g}_{\alpha\beta}$ has the same form, with $e^{\tilde{\nu}}$ and $e^{\tilde{\zeta}}$, related to the Einstein metric functions through $\tilde{\nu} = \nu + 2\phi$, $\tilde{\zeta} = \zeta - 2\phi$. Isotropy implies $\phi = \phi(r)$. Assuming an ideal pressureless matter fluid, $\tilde{T}_{\alpha\beta} = \tilde{\rho}\tilde{u}_\alpha\tilde{u}_\beta$. Motivated by a homogeneous and isotropic cosmology, the vector field is considered time-like. The normalisation condition imposed by the Lagrange multiplier in Eq. (1) gives $U^\alpha = (e^{-\nu/2}, 0, 0, 0)$.

Previous attempts [10, 11, 12] at lensing analysis in TeVeS have remained non-relativistic, considering only the effect of adding a scalar potential to the standard Newtonian potential. Hence they are insensitive to any unique features of TeVeS as a fully relativistic field theory. The analysis given here is used to solve (numerically) the TeVeS equations of motion and obtain explicit expressions for the physical metric quantities $\tilde{\nu}$, $\tilde{\zeta}$ and thus derive the modified Birkhoff's theorem for the TeVeS theory. These functions are used to obtain the deflection angle and find the lensing mass, which is then compared to the stellar mass content. Assuming a mass density profile $m_s(< r)$, within a radial distance r , leads to a system of differential equations to determine ζ and ν . A transformation to $\tilde{\nu}$, $\tilde{\zeta}$ then gives the *physical* metric. The (tt) and $(\theta\theta)$ differential equations are:

$$\begin{aligned} \zeta'' + \frac{(\zeta')^2}{4} + \frac{2\zeta'}{r} + e^\zeta \Lambda = & -\frac{kG^2 m_s^2 e^{-(\nu+\zeta)}}{4\pi\mu(y) r^4} \\ & - \frac{2\pi\mu^2(y)}{l^2 k^2} F(\mu) e^\zeta - K \left[\frac{(\nu')^2}{8} + \frac{\nu''}{2} + \frac{\nu'\zeta'}{4} + \frac{\nu'}{r} \right] \\ & - 8\pi G \tilde{\rho} e^{\zeta-2\phi}, \\ \frac{(\nu' + \zeta')}{2r} + \frac{(\nu')^2}{4} + \frac{\zeta'' + \nu''}{2} + e^\zeta \Lambda = & -\frac{kG^2 m_s^2 e^{-(\nu+\zeta)}}{4\pi\mu(y) r^4} - \frac{2\pi\mu^2(y)}{l^2 k^2} F(\mu) e^\zeta + \frac{K}{8} \nu'^2; \end{aligned} \quad (3)$$

where the prime denotes derivatives with respect to r and the mass profile is $m_s(< r) = 4\pi \int_0^r \tilde{\rho} \exp\left(\frac{\nu}{2} + \frac{3\zeta}{2} - 2\phi r^2\right) dr$. The deflection angle reads [2]:

$$\Delta\phi = 2 \int_0^\infty \frac{1}{r} \left(e^{\tilde{\zeta}-\tilde{\nu}} \frac{r^2}{b^2} - 1 \right)^{-1/2} dr - \pi; \quad (4)$$

$b^2 = e^{\tilde{\zeta}(r_0)-\tilde{\nu}(r_0)} r_0^2$ is the observable impact parameter and r_0 is the point of closest approach for the light ray.

The Navarro-Frenk-White (NFW) profile [9] is used, which in Schwarzschild radial coordinates $\hat{r}$, with $\hat{r} = e^{\tilde{\zeta}/2} r$, reads $M(< \hat{r}) = M \left[\ln\left(1 + \frac{\mathcal{C}\hat{r}}{r_{\text{vir}}}\right) - \frac{\mathcal{C}\hat{r}}{r_{\text{vir}} + \mathcal{C}\hat{r}} \right] \left[\ln(1 + \mathcal{C}) - \frac{\mathcal{C}}{1+\mathcal{C}} \right]^{-1}$, where the concentration $\mathcal{C}$ is $\mathcal{C} \sim 10$ and M is the total mass

of the galaxy contained within the virial radius, r_{vir} ; it also specifies the density profile.

Making the approximation $m_s(< r) \approx M(< r)$, shown [2] to be correct to leading order of r , Eq. (3) is numerically solved and through a transformation to the physical metric, the deflection angle Eq. (4) is obtained.

A specification for the function μ (see, Eq. 2) is required. Previous lensing studies [11, 12] on the non-relativistic scalar potential approach to TeVeS, adopted a model for $\mu(y)$ given in Ref. [2]. It was however noted [14, 15] that when this choice is converted into its MONDian equivalent, rotation curve data are not well fitted. It was hence proposed to consider [16] the MOND function which best fits the rotation curve data and then convert it into its TeVeS $\mu(y)$ analogue. Any intermediate choice for $\mu(y)$ can be parametrised by α . In what follows the results from testing this parametrised $\mu(y)$ in the fully relativistic treatment of lensing in TeVeS are presented in order to see if TeVeS can both fit rotation curve data and lensing data for a single choice of $\mu(y)$. For TeVeS $\mu(y)$, $F(\mu)$ and MOND $f(x)$ we use:

$$\begin{aligned} \mu(y) &= \frac{\sqrt{y/3}}{1 - \frac{4\pi\alpha}{k}\sqrt{y/3}} \\ F(\mu) &= \frac{6k^3}{(4\pi\alpha)^3 \mu^2} \left[\ln\left(\frac{4\pi\alpha\mu}{k} + 1\right)^2 + \frac{1}{1 + \frac{4\pi\alpha\mu}{k}} - \frac{4\pi\alpha\mu}{k} \right] \\ f(x) &= \frac{2x}{1 + (2 - \alpha)x + \sqrt{(1 - \alpha x)^2 + 4x}}, \end{aligned} \quad (5)$$

where the parameter range is $0 < \alpha \leq 1$ [16]. The $\alpha = 0$ case for $\mu(y)$ gives the weak and intermediate gravity limit of $\mu(y)$ and $F(\mu)$ is taken from its explicit form [2]. The $\alpha = 1$ gives the function which better fits rotation curve data. Since the functions increase monotonically, the analysis could be confined to the extremes of the parameter space, i.e. the $\alpha = 0$ and $\alpha = 1$ cases. The TeVeS parameters are [10] $k = 0.01$, $K = 0.01$, $\ell = \sqrt{k\tilde{b}}/(4\pi\Xi a_0)$, $\phi_c = 0.001$, $\Xi = 1 + K/2 - 2\phi_c$.

The cosmological model used is $(\Omega_m, \Omega_\Lambda, \Omega_k) = (0.3, 0.7, 0)$, though the effect of other choices is also considered. The parameter $\tilde{b}$ is fixed from the limit of the $y(\mu)$ function as μ becomes $\ll 1$, $y(\mu) \approx \tilde{b}\mu^2$. For our parametrised choice of $\mu(y)$ $\tilde{b} = 3$, specifying then l . The deflection angle for a model system is then calculated with this choice of parameters. The results for the deflection angle in TeVeS are compared to the deflection angle resulting from following the scalar potential method employed in Refs. [10, 11, 12], as well as GR and MOND. The deflection angle results are shown in Fig. 1.

To calculate a possible dark matter component, a sample of double lensing systems from the Castles database is analysed and the mass of the lensing galaxy in GR, MOND, and TeVeS is calculated. By comparing the mass from lensing to the stellar mass content calculated from a comparison of photometry and stellar population synthe-




FIG. 1: Deflection angle curves for a NFW profile system ($\mathcal{C} = 10$). The parameters of the system are $D_l = 1000$ Mpc (the luminosity distance of the lens), $M = 10^{11} M_\odot$, $k = 0.01$, $K = 0.01$, $\phi_c = 0.001$ and $r_{vir} = 15$ Kpc. *Left panel:* $\alpha = 0$ case. *Right panel:* $\alpha = 1$ case. The deflection angles decrease in MOND and TeVeS as α increases.

sis using a Chabrier Initial Mass Function (IMF), as in Ref. [13], the mass deficit which belonged to the “dark” sector is found. Note that while the authors of Ref. [6] claim that the stellar mass estimates are sensitive to the IMF used, it has been argued in Ref. [7] that the main competitor of the Chabrier IMF, the Salpeter IMF, appears to fit the data worse. Other realistic IMF choices differ from the Chabrier by an insignificant factor [7]. These arguments support the validity of the method employed here. Finally, inverse ray tracing is used to calculate the mass [4]. The mass estimates we obtain this way are shown in Table 1.

Lens	GR	MOND		TeVSeS		M _{STAR} Ref. [13]
		$\alpha = 0$	$\alpha = 1$	$\alpha = 0$	$\alpha = 1$	
HS0818+1227	34.9	23.9	27.6	19.1	24.4	$16.2^{21.2}_{12.6}$
FBQ0951+2635	3.1	2.3	2.6	1.9	2.3	$1.1^{2.1}_{0.5}$
BRI0952-0115	3.6	2.4	2.8	1.9	2.4	$3.5^{4.0}_{2.7}$
HE1104-1805	86.1	59.0	68.5	47.1	60.4	$22.8^{51.2}_{12.7}$
LBQS1009-0252	15.2	10.7	12.2	8.7	10.9	$5.5^{7.9}_{4.2}$
HE2149-2745	11.2	7.8	9.0	6.3	8.0	$4.6^{6.7}_{3.6}$

TABLE I: Mass estimates (in $10^{10} M_\odot$ units) for Λ CDM cosmology: $(\Omega_m, \Omega_\Lambda, \Omega_k) = (0.3, 0.7, 0)$, $k = K = 0.01$, and a NFW halo profile with $\mathcal{C} = 10$. Two different cases ($\alpha = 0, 1$) for $\mu(y)$ parametrisation are considered.

Figure 2 compares the mass estimates between GR, MOND and TeVeS for the two cases of $\mu(y)$, with the mass difference being given as a function of the mass calculated using GR (left panels) and R_{lens}/R_e (right panels), where R_{lens} is the distance out to which our mass estimates and the stellar mass estimates are calculated and R_e is the half light radius, both given in Ref. [13]. The top two panels are for $\alpha = 0$, the bottom two are for $\alpha = 1$. Comparing the lensing masses against the stel-

lar masses (excluding one outlier), it is found that even in TeVeS on average the dark matter content is 48.5% when $\alpha = 0$, and 34.3% when $\alpha = 1$. The outlier lensing system BRI0952-0115 shows a mass overshoot of $\approx 80\%$ in TeVeS, i.e. the lensing mass in TeVeS is less than the stellar mass content. It is possible that this system is affected by some unknown lens environment effects such as an unseen cluster mass contribution as has been suggested of other lenses [11], though there is no data to conclude this at present. Overall the analysis shows that TeVeS finds it hard to explain the lensing observed in these systems using only the stellar content of the galaxies, a conclusion which stands in contrast to that given in Refs. [11, 12] despite the two methods predicting similar deflection angles (see, Fig.1). This is a problem for TeVeS, which will be made more explicit later on in this letter, when the analysis of varying the free parameters will be performed.




FIG. 2: A comparison of the need for dark matter for the lenses in GR, MOND and TeVeS, for $\alpha = 0$ (top panels) and $\alpha = 1$ (bottom panels). The right hand panels give the dark matter requirements as a function of the ratio R_{lens}/R_e . Any dark matter needed in TeVeS is especially significant for those masses which probe further out, where the modification to gravity is larger.

The results show that the case $\alpha = 1$, which is a choice specifically adopted to fit the rotation curve data, requires an even larger amount of dark matter. Thus, attempts to fit TeVeS to a no dark matter scenario using the freedom in the parameters of the class of μ functions used here and in the literature so far, would imply that the theory fits poorly the rotation curves data, which was the original motivation for modifying the gravitational behaviour.

The effect of different cosmological parameters has also been examined to see how they alter the results for the

$\alpha = 0$ case, corresponding to the minimum amount of required dark matter. It is found that the lensing results are insensitive to the precise cosmological parameters, which is not surprising, since the observational constraints mostly impose limits on the luminosity and angular distance scales [4]. For completeness we state the results, all of which point towards fluctuations in the amount of dark matter well within the error limits. In particular, for the case [12] $(\Omega_m, \Omega_\Lambda, \Omega_k) = (0.03, 0.46, 0.51)$, averaging over the six galaxies, we find a 5.7% increase in the amount of dark matter required. For the case $(\Omega_m, \Omega_\Lambda, \Omega_k) = (0.23, 0.78, 0)$, considered in Ref. [8], to fit the Cosmic Microwave Background data with TeVeS, one finds a corresponding average decrease by 2.2 %, while for $\phi_c = 0.01$, one finds an average decrease by 1.1% .

Finally, the k , K parameters are varied independently to check on the robustness of our claims. In particular, we examine the lensing system HE1104-1805, which within the TeVeS approach (with $\alpha = 0$) requires the largest amount of dark matter. We find that the variation of the parameter k has considerably smaller effects than that of K . This supports the dominant rôle played by the vector field in TeVeS. A similar result has also been pointed out in Ref. [17], but from a different perspective. Within the allowed parameter space $1 \leq K \leq 10^{-5}$, $1 \leq k \leq 10^{-5}$, implied by rotation curve data and solar system tests of gravity, the amount of dark matter required in the system HE1104-1805 is negligible only for values of K higher than 0.1, which however is excluded by gravitational measurements at solar-system scales.




FIG. 3: The dark matter content of the lenses for $\alpha = -1$ in the parametrisation of $f(x)$ in MOND and $\mu(y)$ in TeVeS.

We also considered other values of α , outside the region $\alpha \in [0, 1]$, considered in the literature so far. We found that the $\alpha = -1$ case appears not to require substantial amount of dark matter to explain the gravitational lensing, and hence it could provide an example of an alternative to the dark matter scenario. However, such a model is in conflict with data from rotation curves of

galaxies.

In conclusion, the above results show a soft dependence on the free parameters of TeVeS and the cosmological model adopted, but a rather strong one on the form of the $\mu(y)$ function. Our analysis in this letter points towards the fact that TeVeS, at least within the class of models considered so far in the literature, cannot survive both gravitational lensing and rotation curve tests. However, we cannot yet exclude completely the possibility, admittedly remote, that a class of μ functions, or more complicated equivalents thereof, can be found such that alternative to dark matter scenarios are at play.

It is a pleasure to thank Ignacio Ferreras for discussions. The work of N.E.M. and M.S. is partially supported by the European Union through the Marie Curie Research and Training Network UniverseNet (MRTN-CN-2006-035863), while that of M.F.Y. is supported by an E.P.S.R.C. (UK) studentship.

* Electronic address: Nikolaos.Mavromatos@kcl.ac.uk

† Electronic address: Mairi.Sakellariadou@kcl.ac.uk

‡ Electronic address: Muhammad.Yusaf@kcl.ac.uk

- [1] M. Milgrom, *Astrophys. J.* **270** (1983) 365.
- [2] J. D. Bekenstein, *Phys. Rev. D* **70** (2004) 083509; R. H. Sanders, *Astrophys. J.* **473** (1996) 117.
- [3] N. E. Mavromatos and M. Sakellariadou, *Phys. Lett. B* **652** (2007) 97.
- [4] I. Ferreras, M. Sakellariadou and M. F. Yusaf, *Phys. Rev. Lett.* **100** (2008) 031302
- [5] R. H. Sanders, *Mon. Not. R. Astron. Soc.* **380** (2007) 331.
- [6] R. H. Sanders, D. D. Land [arXiv:astro-ph/0803.0468v2].
- [7] I. Ferreras, P. Saha, S. Burles, *Mon. Not. Roy. Astron. Soc.* **383** (2008) 857.
- [8] C. Skordis, D. F. Mota, P. G. Ferreira and C. Boehm, *Phys. Rev. Lett.* **96** (2006) 011301.
- [9] J. F. Navarro, C. S. Frenk, S. D. M. White, *Astrophys. J.* **493** (1996) 563.
- [10] M. Feix, C. Fedeli and M. Bartelmann, *Astron. Astrophys.* **480** (2008) 313.
- [11] M. Feix, H. Shan, B. Famaey, H. Zhao [arXiv:0804.2668v1]
- [12] H. Zhao, D. J. Bacon, A. N. Taylor and K. Horne, *Mon. Not. R. Astron. Soc.* **368** (2006) 171.
- [13] I. Ferreras, P. Saha and L. L. R. Williams, *Astrophys. J.* **623** (2005) L5.
- [14] H. Zhao, B. Famaey, *Mon. Not. Roy. Astron. Soc.* **368** (2006) 171.
- [15] B. Famaey, J. Binney, *Mon. Not. Roy. Astron. Soc.* **363** (2005) 603.
- [16] G. W. Angus, B. Famaey, H. Zhao, *Mon. Not. Roy. Astron. Soc.* **371** (2006) 138.
- [17] S. Dodelson and M. Liguori, *Phys. Rev. Lett.* **97** (2006) 231301.